

CA

A Cancer Journal for Clinicians

H.G. Wells on Cancer

Howard W. Kays

CA Cancer J Clin 1982;32:63-64

DOI: 10.3322/canjclin.32.1.63-b

This information is current as of March 16, 2010

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://caonline.amcancersoc.org>

To subscribe to the print issue of *CA: A Cancer Journal for Clinicians*, go to (US individuals only): <http://caonline.amcancersoc.org/subscriptions/>

CA: A Cancer Journal for Clinicians is published six times per year for the American Cancer Society by Wiley-Blackwell. A bimonthly publication, it has been published continuously since November 1950. *CA* is owned, published, and trademarked by the American Cancer Society, 250 Williams Street NW, Atlanta GA 30303. (©American Cancer Society, Inc.) All rights reserved. Print ISSN: 0007-9235. Online ISSN: 1542-4863.



Reference

1. Bittner JJ: Possible method of transmission of susceptibility to breast cancer in mice. *Am J Cancer* 39:104–113, 1940.

Author's Reply:

Epidemiologic studies have established that breast-feeding as practiced in the United States bears little relationship to breast cancer (see for instance Wynder EL, Bross IJ, Hirayama T: A study of the epidemiology of cancer of the breast. *Cancer* 13:559–601, 1960). Dr. Pollak's conjectures may have surface plausibility, but on further reflection I do not see how recent small presumed changes in such an unimportant factor as breast-feeding can be given much credence for explaining the trends in breast cancer incidence in younger women.

LaSalle D. Leffall, Jr., M.D.
Professor and Chairman
Department of Surgery
Howard University Hospital
Washington, D.C.

Postmastectomy Lymphedema

To the Editor:

I read with interest your Classics in Oncology article, "Lymphangiosarcoma in Postmastectomy Lymphedema" (*Ca* 31:284–299, 1981). I feel it is imperative to respond to this article because much is said about the syndrome of lymphangiosarcoma occurring after long-term lymphedema of the arm, but little or nothing constructive is said regarding prevention.

If one can conclude that postmastectomy lymphedema of the arm is in fact a cause of late lymphangiosarcoma (or other sarcomas) of the arm, then prevention of lymphedema should clearly be a method of prevention. In support of this supposition I am unaware of any postmastectomy patients developing angiosarcoma of the arm in the absence of lymphedema.

If postmastectomy lymphangiosarcoma does not occur in the absence of lymphedema, then the prevention of lymphedema should in fact be essential in the prevention of lymphangiosarcoma, but also in the prevention of painful, distressing, recurrent lymphangitis and cellulitis.

My point in writing this letter is to stress the importance of preventing long-term lymphedema by careful follow-up of these patients with the judicious use of diuretics and the application of compressive-type elastic sleeves as appropriate and/or the use of intermittent pressure devices. If these measures are not instituted promptly following the mastectomy, collagen deposition occurs in the interstitial tissues and the swelling becomes an irrevocable situation. Other factors that seem from the clinical standpoint to be important in the causation of this chronic lymphedema are pre- or postoperative irradiation of the axilla and possibly overzealous surgical dissection high in the axilla.

I feel that if these preventative measures are kept in mind, many of these chronic lymphedema problems can be avoided in the future.

Gordon R. Held, M.D.
Associate Professor of Surgery
University of South Dakota
Royal C. Johnson Veterans
Memorial Hospital
Sioux Falls, South Dakota

H.G. Wells on Cancer

To the Editor:

In the September/October Feedback section (*Ca* 31:320, 1981), Dr. Cohen misinterpreted the quotation of H.G. Wells: "... the motive that will conquer cancer will not be pity nor horror; it will be curiosity to know how and why."

H.G. Wells did *not* mean that scientists are doubtful humanitarians not wishing to help people with killing diseases.

He *did* mean that cancer will not be considered a dreaded and horrible disease unlike any other. Previous articles in *Ca* have shown that only a few years ago cancer patients were not even told their diagnosis, because cancer was associated with a horrible, wasting, painful death. Since people were afraid to be reminded of their own mortality, cancer patients were ostracized like lepers.

In 1927, H.G. Wells correctly predicted the decreasing fear of cancer, as scientific curiosity discovers better treatment and prevention.

Howard W. Kays, M.D.
A.C.S. Clinical Oncology Fellow
Department of Radiation Oncology
Indiana University Hospitals
Indianapolis, Indiana